

**PORTLAND VETERANS AFFAIRS MEDICAL CENTER
INSTITUTIONAL ANIMAL CARE AND USE COMMITTEE (PVAMC IACUC)
ANIMAL CARE AND USE GUIDELINES**

CHECK ONE:

- I agree to comply with the following guidelines.
- I have attached written justification for deviation from these guidelines.

Principal Investigator

Date

GUIDELINES ON DEATH AS AN ENDPOINT

Background. The Office for Protection from Research Risks (OPRR) *Institutional Animal Care and Use Committee Guidebook (NIH Publication No. 92-3415)* states that “Endpoints other than death must always be considered and should be used whenever the research objective makes it possible.”

Guidelines.

1. The PVAMC IACUC discourages the use of death as an endpoint, and instead encourages the use of alternative endpoints. Examples of alternative endpoints are given below (endpoint utilized determined by type of experimental application, e.g., tumor study, toxicity testing, surgical procedure, etc.):
 - a. Tumor load > 10% of normal body weight.
 - b. Weight loss of > 20% of normal body weight (due to anorexia, physical inability to obtain food and/or water, or severe diarrhea or vomiting).
 - c. CNS signs such as circling, blindness, convulsions.
 - d. Hematologic parameters indicative of end-stage organ failure.
2. Animals should be humanely euthanized at the time the alternative endpoint is reached, or if they demonstrate signs of being moribund. *Moribund* is defined as “in a dying state” (Dorland’s Medical Dictionary, 26th Ed., Philadelphia: W.B. Saunders, 1981). Animals are considered to be moribund if they manifest any of the following clinical signs and recovery is not expected (e.g., recovery would be expected with anesthesia):
 - a. Inability to maintain an upright position.
 - b. Prolonged (greater than 48 hours) physical inability to obtain food and/or water.
 - c. Prolonged (greater than 48 hours) anorexia and/or clinical dehydration.
 - d. Uncontrollable diarrhea, vomiting or constipation.
 - e. Agonal breathing and cyanosis.
 - f. Unconsciousness with no response to external stimuli (e.g., toe-pinch withdrawal test).
3. If death as an endpoint must be used (i.e., killing a moribund animal would invalidate the study) the following stipulations must be met:
 - a. Scientific justification for using death as an endpoint must be provided in writing as part of the animal care protocol and must be approved by the PVAMC IACUC.
 - b. Moribund animals must be monitored at minimum twice daily (in the early morning and late afternoon, including weekends and holidays).
 - c. Written records must be made of all monitoring sessions indicating the time and date of the observation, the person performing the observation, and any findings

(such as number of animals demonstrating clinically abnormal behavior, number of animals found dead, etc.). These records must be kept on file and made available to the VMU personnel and/or IACUC on request.

- d. Moribund animals must be removed from group housing and housed individually with easy access to food and water.
- e. The minimum number of animals necessary to achieve statistical significance must be used.
- f. Drugs or techniques to alleviate pain or distress preceding death must be used unless they would interfere with the scientific objectives of the study.
- g. Proposals foregoing the use of anesthetics, analgesics or tranquilizing drugs must be extensively justified in writing as part of the animal care protocol and approved by the PVAMC IACUC. Additionally, proposals which utilize death as an endpoint and which forego the use of anesthetics, analgesics or tranquilizing drugs to alleviate pain and distress in experimental animals will be assigned the highest pain level category, "E," on all protocol forms and regulatory papers.